

Paper Number: 4756

Reconstructing the plumbing system of Krakatau volcano

Troll, V.R.^{1,2}, Deegan, F.M.¹, Dahren, B.¹, Jolis, E.M.¹, Blythe, L.S.¹, Harris, C.³, Berg, S.E.¹, Hilton, D.R.⁴, and Freda, C.²

¹ CEMPEG, Uppsala University, Uppsala, Sweden. Email: Valentin.Troll@geo.uu.se

² Istituto Nazionale di Geofisica e Vulcanologia (INGV), Rome, Italy, Rome, Italy.

³ University of Cape Town, Cape Town, South Africa.

⁴ Scripps Institution of Oceanography, San Diego, CA, United States.

Crustal contamination of ascending arc magmas is generally thought to be significant at lower- to mid-crustal magma storage levels where magmas inherit their chemical and isotopic character by blending, assimilation and differentiation [1]. Anak Krakatau, like many other volcanoes, erupts shallow-level crustal xenoliths [2], indicating a potential role for upper crustal modification and hence late-stage changes to magma rheology and thus eruptive behaviour. Distinguishing deep vs. shallow crustal assimilation processes at Krakatau, and elsewhere, is therefore crucial to understand and assess pre-eruptive magmatic conditions and their associated hazard potential. Here we report on a multi-disciplinary approach to unravel the crustal plumbing system of the persistently-active and dominantly explosive Anak Krakatau volcano [2, 3]. We employ rock-, mineral- and gas-isotope geochemistry and link these results with seismic tomography [4]. We show that pyroxene crystals formed at mid- and lower-crustal levels (9-11 km) and carry almost mantle-like isotope signatures (O, Sr, Nd, He), while feldspar crystals formed dominantly at shallow levels (< 5km) and display unequivocal isotopic evidence for late stage contamination (O, Sr, Nd). Coupled with tomographic evidence, the petrological and geochemical data place a significant element of magma-crust interaction (and hence magma storage) into the uppermost, sediment-rich crust beneath the volcano. Magma – sediment interaction in the uppermost crust therefore offers a likely explanation for the compositional variations in recent Krakatau magmas and most probably provides extra impetus to increased explosivity at Anak Krakatau.

References:

- [1] Annen et al. (2006) J Petrol 47:505-539
- [2] Gardner et al. (2013) J Petrol 54:149-182
- [3] Dahren et al. (2012) Contrib Mineral Petrol 163:631-651
- [4] Jaxybulatov et al. (2011) J Volcanol Geotherm Res 206:96-105.

